

BAIRE REFLECTION

STEVO TODORCEVIC AND STUART ZOBLE

ABSTRACT. We study reflection principles involving nonmeager sets and the Baire Property which are consequences of the generic supercompactness of ω_2 , such as the principle asserting that any point countable Baire space has a stationary set of closed subspaces of weight ω_1 which are also Baire spaces. These principles entail the analogous principles of stationary reflection but are incompatible with forcing axioms. Assuming MM , there is a Baire metric space in which a club of closed subspaces of weight ω_1 are meager in themselves. Unlike stronger forms of Game Reflection, these reflection principles do not decide CH , though they do give ω_2 as an upper bound for the size of the continuum.

INTRODUCTION AND BASIC THEORY

For a set A of ω -sequences of ordinals and a set of ordinals H the game $G(A, H)$ has two players who alternate playing ordinals from H . Player II wins if the cooperative play belongs to A and loses otherwise. A weak version of the Game Reflection Principle defined and studied in [9], which we denote $GRP_\omega(\theta)$ for an uncountable cardinal θ , asserts that for every $A \subset \theta^\omega$, player II has a winning strategy in $G(A, \theta)$ if and only if II has a winning strategy in $G(A, H)$ for an ω_1 -club of $H \in [\theta]^{\omega_1}$. Here an ω_1 -club means the set of H closed under some function $f : \theta^{<\omega} \rightarrow \theta$.¹ The weakened reflection principle obtained by requiring each player to play finite sequences, rather than single ordinals (producing an element of θ^ω by concatenation of the plays), is immediately equivalent to the Weak Baire Reflection Principle below.

Definition 0.1. $BRP^w(\theta)$ asserts that any $A \subset \theta^\omega$ is meager if and only if $A \cap H^\omega$ is meager in H^ω for an ω_1 -club of $H \in [\theta]^{\omega_1}$.

A stronger version requires reflection of a failure of the Baire Property.

Definition 0.2. $BRP(\theta)$ asserts that any $A \subset \theta^\omega$ has the Baire Property in θ^ω if and only if $A \cap H^\omega$ has the Baire Property in H^ω for an ω_1 -club of $H \in [\theta]^{\omega_1}$.

BRP^w and BRP will denote the global versions of these principles. Before deducing BRP from Game Reflection we discuss how to expand the scope of these principles to spaces other than the Generalized Baire Spaces (spaces of the form θ^ω). Recall that the weight of a space X is the minimum cardinality of a base for

Received by the editors March 10, 2006.

2000 *Mathematics Subject Classification.* Primary 03E55; Secondary 03E50.

Key words and phrases. Baire Property, Game Reflection, Martin's Maximum.

¹In [9] the less restrictive notion of a set closed under ω_1 -length increasing unions and cofinal in the \subseteq -ordering is used.

the topology of X . The point weight of a space X is defined to be the minimum cardinal κ such that X has a base \mathcal{B} which satisfies

$$|\{b \in \mathcal{B} \mid x \in b\}| \leq \kappa$$

for every $x \in X$. A point countable space is a space with point weight ω . For a space X with base \mathcal{B} and a point $x \in X$ let $\mathcal{B}_x = \{b \in \mathcal{B} \mid x \in b\}$. For a set $B \subseteq \mathcal{B}$ let $X_B = \{x \in X \mid \mathcal{B}_x \subseteq B\}$. Note that X_B is always a closed subspace of X . By an ω_1 -club of subspaces of X of weight ω_1 we will mean a set of the form $\{X_B \mid B \in C\}$ for some ω_1 -club $C \subseteq [\mathcal{B}]^{\omega_1}$. We may thus speak of $BRP(\theta)$ ($BRP^\omega(\theta)$) for any class of spaces: for any member space X with weight less than or equal to θ and $A \subseteq X$, A has the Baire Property (is meager) in X if and only if $A \cap Y$ has the Baire Property (is meager) in Y for an ω_1 -club of subspaces $Y \subseteq X$ of weight ω_1 . It is easy to see that any ω_1 -club of subspaces of λ^ω contains a club of the form $\{Y^\omega \mid Y \in C\}$ for some ω_1 -club $C \subseteq [\lambda]^{\omega_1}$, so that Baire Reflection as stated in Definitions 0.1 and 0.2 is equivalent to Baire Reflection for the class of generalized Baire spaces.

Theorem 0.3. *$GRP_\omega(\theta)$ implies $BRP(\theta)$ for completely regular spaces of point weight less than or equal to ω_1 .*

Proof. Fix such a space X of weight θ . Let $\{N_\alpha \mid \alpha \in \theta\}$ enumerate such a base \mathcal{B} for the topology on X with no repetitions, and let $A \subseteq X$ have the reflection property as witnessed by a club $C \subseteq [\mathcal{B}]^{\omega_1}$. A key point is that C can be refined to a club $C_0 \subset C$ with the property that $b \cap X_B \neq \emptyset$ for each $b \in B \in C$. Simply take C_0 to be the set of $\mathcal{B} \cap H$ in C for an elementary submodel $H \prec H(\kappa)$ with $\omega_1 \cup \{X, \mathcal{B}\} \subset H$. We describe a game $G(A^*, \theta)$ that simulates the relevant Banach-Mazur games with player II as the winner. A^* is the set of sequences $(\alpha_i) \in \theta^\omega$ such that

- (1) $N_{\alpha_{i+1}} \subseteq N_{\alpha_i}$ for every $i \in \omega$ and the containment is proper for every i which is nonzero,
- (2) if α_0 is equal to α_1 , then $\bigcap_{i < \omega} N_{\alpha_i} \cap A = \emptyset$,
- (3) if α_0 is not equal to α_1 , then $\bigcap_{i < \omega} N_{\alpha_i} \subseteq A$.

Now consider an $H \in [C_0]^{\omega_1}$ in the club C_0 . H is a base for the subspace topology on X_H . Thus the relevant Banach-Mazur games for $A \cap X_H$ are determined. These strategies give rise to a winning strategy for II in the game $G(A^*, H^*)$ in the obvious way where H^* is the set of α with $N_\alpha \in H$. Thus II has a winning strategy in $G(A^*, \theta)$ and this in turn gives rise to the Banach-Mazur strategies which show that A has the Baire Property in X . \square

For an infinite cardinal κ let Γ_κ^{UB} denote the pointclass of κ -Universally Baire sets of reals.

Corollary 0.4. *$GRP_\omega(\kappa)$ implies $\Gamma_{\omega_1}^{UB} = \Gamma_\kappa^{UB}$. If κ is a large enough cardinal (for example a huge cardinal) and $G \subset Col(\omega_1, < \kappa)$ is V -generic, then in $V[G]$ every set of reals in $\Gamma_{\omega_1}^{UB}$ is determined.*

Proof. Game Reflection and hence BRP holds in $V[G]$ by Theorem 0.3 above and Cor. 18 of [9]. The conclusion follows as Universally Baire sets of reals are determined in the presence of two Woodin cardinals (see [4]). \square

We now show that ostensibly stronger versions of BRP and BRP^ω are equivalent to the versions stated in Definitions 0.1 and 0.2. Recall that the cellularity of a space

is the minimum cardinal λ such that any pairwise disjoint collection of nonempty open sets has size less than λ . In a metric space, λ is the maximum cardinality of such an antichain. This is because it suffices to consider families of basic open sets, and metric spaces have σ -discrete bases by Stone's Theorem (see [2]). Let us call a space X a Baire space, or everywhere Baire, if the intersection of any family $\{D_n \mid n \in \omega\}$ of dense open sets is dense. If there is a sequence of open dense sets with empty intersection, then we say that X is meager in itself.

Theorem 0.5. *For any uncountable cardinal θ the principle $BRP^\omega(\theta)$ is equivalent to its version for metric spaces.*

Proof. Assume $BRP^\omega(\theta)$ and let M be a metric space of weight less than or equal to θ . Let \mathcal{B} be a σ -discrete base for M consisting of ϵ -balls. Suppose $A \subseteq M$ has the reflection property, that is, suppose there is an ω_1 -club $C \subseteq [\mathcal{B}]^{\omega_1}$ such that $A \cap Y_B$ is meager in Y_B for any $B \in C$. Fix a maximal pairwise disjoint family \mathcal{U} of elements of \mathcal{B} with the property that every $U \in \mathcal{U}$ is everywhere λ_U cellular for some cardinal λ_U . By everywhere λ_U cellular we mean that the cellularity of every nonempty open $V \subseteq U$ is also λ_U . \mathcal{U} can be obtained by collecting the minimal elements of the well-founded relation on $\mathcal{B} \setminus \{\emptyset\}$ defined by $U < V$ if $U \subseteq V$ and $c(U) < c(V)$. Clearly the union of \mathcal{U} is dense in M . It suffices to show that $A \cap U$ is meager in U for every $U \in \mathcal{U}$. If $\lambda_U \leq \omega_1$, then there is $B \in C$ which contains $\{b \in \mathcal{B} \mid b \cap U \neq \emptyset\}$ so that $U \subseteq X_B$. It follows that $A \cap U$ is meager in U in this case. Thus we concentrate on $U \in \mathcal{U}$ with $\lambda_U > \omega_1$. Let \mathcal{B}_U denote the set of $b \in \mathcal{B}$ such that $b \subseteq U$. Thus \mathcal{B}_U is a base for the subspace topology on U . Note that B_U has size λ_U . Let $C_U = \{B \cap \mathcal{B}_U \mid B \in C\}$. C_U is an ω_1 -club in $[\mathcal{B}_U]^{\omega_1}$. Let U^* be the metric completion of U . We show that there is a dense G_δ set $G \subseteq U^*$ and a homeomorphism $\pi : G \rightarrow \lambda_U^\omega$. Let $\lambda_U = \lambda$. For a finite sequence $s \in \lambda^{<\omega}$ let $lh(s)$ denote the length of s . For $b \in \mathcal{B}_U$ let b^* be the extension to U^* . The set $\mathcal{B}_U^* = \{b^* \mid b \in \mathcal{B}_U\}$ is a base for U^* . By induction there is a map $\phi : \lambda^{<\omega} \rightarrow \mathcal{B}_U^*$ such that $\phi(\emptyset) = U$ and

- (1) $lh(s) = n$ implies $\phi(s)$ has radius less than $\frac{1}{n}$,
- (2) $s \subset t$ implies $\phi(t) \subset \phi(s)$,
- (3) $\bigcup \phi[\lambda^n]$ is dense in U ,
- (4) s not equal to t implies $cl_{U^*}(\phi(s)^*) \cap cl_{U^*}(\phi(t)^*)$ is empty.

Thus G defined by

$$G = \bigcup_{f \in \lambda^\omega} \bigcap_{n \in \omega} \phi(f \upharpoonright n)^* = \bigcap_{n \in \omega} \bigcup_{s \in \lambda^n} \phi(s)^*$$

is the desired G_δ with $\pi^{-1}(f)$ defined to be the unique element of $\bigcap_{n \in \omega} \phi(f \upharpoonright n)$. The map π acts on elements of \mathcal{B}_U^* and subsets of \mathcal{B}_U^* . Let $C_U^* = \{B^* \mid B \in C\}$ where $B^* = \{b^* \mid b \in B\}$. Thus C_U^* is a club in $[\mathcal{B}_U^*]^{\omega_1}$. Let C_U^{**} be the projection of C_U^* on $\{\phi(s)^* \mid s \in \lambda^{<\omega}\} = \mathcal{B}_U^{**}$. Thus $\pi[C_U^{**}]$ is club in $[\lambda^{<\omega}]^{\omega_1}$ (identifying basic open sets in λ^ω with elements of $\lambda^{<\omega}$). For $B \in C_U$ we have

$$X_{B^*}^{U^*} \cap U = X_B^U,$$

where the superscript indicates the parent space, so that A is meager in $X_{B^*}^{U^*}$. Thus

$$X_{B^*}^{U^*} \cap G = X_{B^* \cap \mathcal{B}_U^{**}}^G.$$

It follows that $A \cap G$ is meager in X_B^G for any $B \in C_U^{**}$ so that $\pi[A \cap G]$ is meager in $\pi[X_B^G] = X_{\pi[B]}^{\lambda^\omega}$. $BRP^\omega(\lambda)$ holds as $\lambda \leq \theta$ so $\pi[A \cap G]$ is meager in λ^ω and hence $A \cap G$ is meager in G . It follows that A is meager in U as desired. \square

Theorem 0.6. *For any uncountable cardinal θ , $BRP(\theta)$ is equivalent to its version for complete metric spaces.*

Proof. Assume M is complete metric and $A \subset M$. Again let \mathcal{B} be a σ -discrete basis for M consisting of ϵ -balls, and assume $A \cap Y_B$ has the Baire Property in Y_B for all B in some club $C \subset [\mathcal{B}]^{\omega_1}$. Let $\mathcal{U} \subset \mathcal{B}$ be as in the theorem above. Again, it suffices to show that $A \cap U$ has the Baire Property in U for $U \in \mathcal{U}$ with $\lambda_U > \omega_1$. Since M is already complete, the construction of Theorem 0.5 gives a dense open $G_U \subset U$ which is homeomorphic via a map π to λ_U^ω and it suffices to show that $A \cap G_U$ has the Baire Property in G_U . This is deduced by a similar argument as in the theorem above, using π and our assumption that BRP^w holds in λ_U^ω . \square

We do not know if BRP^w for arbitrary point countable spaces is implied by BRP^w . We also do not know whether BRP for metric spaces or point countable spaces is a stronger principle than BRP . We close this section by defining, in analogy with simultaneous stationary reflection, the Simultaneous Weak Baire Reflection Principle, denoted $BRP_{(2)}^w(\theta)$, to assert that pairs of second category sets $X, Y \subset \theta^\omega$ reflect to second category sets in a stationary set of closed subspaces H^ω of weight ω_1 . It will be shown in the next section that $BRP_{(2)}^w(\theta)$ implies, and is strictly stronger than, $BRP(\theta)$. Again, nothing more is obtained by expanding the scope of this principle to metric spaces. In analogy with diagonal reflection of stationary sets we have the following principle $BRP_d^w(\theta)$ which is yet stronger.

Theorem 0.7. *Assume $GRP_\omega(\theta)$. Suppose X is a completely regular Baire space with a point countable basis. Then a stationary set of closed subspaces $Y \subset X$ of weight ω_1 are also Baire spaces.*

Proof. This is similar to Theorem 0.3 so we omit some detail. Let $\{N_\alpha \mid \alpha < \theta\}$ enumerate a point countable basis \mathcal{B} for X with no repetitions. Assume to the contrary that there is a club $C \subset [\mathcal{B}]^{\omega_1}$ such that Y_B is not a Baire space for $B \in C$. Thus there is a function $f : C \rightarrow \theta$ such that $N_{f(B)} \in B$ and $Y_B \cap N_{f(B)}$ is meager in itself for every $B \in C$. It is easy to see that an equivalent version of $GRP_\omega(\theta)$ is obtained if player II always moves first. Let us assume this version and define $A \subset \theta^\omega$ to be the set of sequences (α_n) such that N_{α_n} is a decreasing chain with empty intersection. For games associated to a $B \in C$, player II wins by using $f(B)$ as a first move and following the winning strategy for “empty” in the Banach-Mazur game for $Y_B \cap N_{f(B)}$. Thus II wins the unrestricted game $G(A, \theta)$ with some first move α , and it follows that N_α is meager in itself, which is a contradiction. \square

The plan for the rest of the paper is as follows. In Section 2, a relationship with stationary reflection is established and used to show that BRP^w does not imply $BRP(\omega_2)$. Throughout, we will strengthen the usual formulation of principles of stationary reflection to assert that reflection occurs stationarily often, where a subset of $[\kappa]^{\omega_1}$ is stationary if it meets every ω_1 -club. These stronger principles follow from Game Reflection as well, though not necessarily from their usual versions (see [3] for more on this issue). One nonstandard principle we use is the following.

Definition 0.8. $WRP_{(2)}^c(\kappa)$ asserts that for any pair of stationary sets $S, T \subset [\kappa]^\omega$ whose union contains a club in $[\kappa]^\omega$, there is a stationary set of $X \in [\kappa]^{\omega_1}$ such that $S \cap [X]^\omega$ and $T \cap [X]^\omega$ are both stationary in $[X]^\omega$.

We will show that $BRP^w(\kappa)$ implies $WRP(\kappa)$ and that $BRP(\kappa)$ implies $WRP_{(2)}^c(\kappa)$ by associating the space $X(S) = \{x \in \kappa^\omega \mid \text{ran}(x) \in S\}$ to a stationary set $S \subseteq [\kappa]^\omega$. Further, we will show that $BRP(\kappa)$ is equivalent to the conjunction of $WRP_{(2)}^c(\kappa)$ and $BRP^w(\kappa)$. We use this connection to produce a model in which BRP^w holds but $BRP(\omega_2)$ does not, namely the model obtained by collapsing a supercompact cardinal to ω_2 and then adding a pair of stationary subsets of $[\omega_2]^\omega$ which do not simultaneously reflect. In Section 2 we show that BRP holds in the model obtained by the Mitchell collapse of a supercompact (see [11]), which shows that BRP does not imply CH , in contrast to König's theorem that $GRP_\omega(\omega_2)$ does imply CH (Thm. 8 of [9]). In the last section of this paper we show that Weak Baire Reflection fails under Martin's Maximum in a variety of ways. For example, assuming MM , we show that there is a Baire metric space of weight ω_2 which has an ω_1 -club of closed subspaces of weight ω_1 which are meager in themselves, and that any set of reals of size ω_1 is ω_1 -Universally Baire but not ω_2 -Universally Baire.

1. RELATIONSHIP WITH STATIONARY REFLECTION

The principle $WRP_{(2)}(\kappa)$ asserts that pairs of stationary sets $S, T \subset [\kappa]^\omega$ reflect simultaneously in the sense that there is a stationary set of $X \in [\kappa]^{\omega_1}$ such that both $S \cap [X]^\omega$ and $T \cap [X]^\omega$ are stationary in $[X]^\omega$. $WRP_{(2)}^c(\omega_2)$ will only apply to sets S, T whose union contains a club.² $WRP_d(\kappa)$ asserts that for any sequence $\{S_\alpha \mid \alpha < \kappa\}$ of stationary subsets of $[\kappa]^\omega$ there is a stationary set of $H \in [\kappa]^{\omega_1}$ such that $S_\alpha \cap [H]^\omega$ is stationary in H^ω for every $\alpha \in H$.

Theorem 1.1. *The following implications hold for any uncountable cardinal κ .*

- (1) $BRP^w(\kappa)$ implies $WRP(\kappa)$.
- (2) $BRP(\kappa)$ implies $WRP_{(2)}^c(\kappa)$.
- (3) $BRP_{(2)}^w(\kappa)$ implies $WRP_{(2)}(\kappa)$.
- (4) $BRP_d^w(\kappa)$ implies $WRP_d(\kappa)$.

Proof. For $S \subset [\kappa]^\omega$ let $X(S) \subset \kappa^\omega$ consist of all sequences x whose range belongs to S . The key point is that $X(S)$ is meager if and only if S is nonstationary. For the reverse direction assume C is a club disjoint from S and let $f : \kappa^{<\omega} \rightarrow \kappa$ be such that $C_f \subset C$. f gives rise to a winning strategy for II in the Banach-Mazur game for $\kappa^\omega \setminus X(S)$. II simply ensures closure of the range of the final play under the function f . For the other direction, assume II wins the Banach-Mazur game. Let C be the set of $\sigma \in [\kappa]^\omega$ such that σ is the range of a play by this strategy. Then C is a club disjoint from S . The preceding holds just as well in any H^ω with H uncountable and so the implication (1) follows. For (2) let S and T be complimentary mod club and both stationary. Assume there is a club $C \subset [\kappa]^{\omega_1}$ such that S, T do not reflect simultaneously at any $H \in C$. Let $X(S) \subset \kappa^\omega$ be the space associated to S and let $H \in C$ be arbitrary. Since $S \cap [H]^\omega$ either

²Paul Larson and James Cummings independently pointed out to the second author that $WRP_{(2)}^c(\omega_2)$ does not imply $WRP_{(2)}(\omega_2)$; see the remarks following 1.7. Note that nothing is lost by requiring that $S \cup T = [\kappa]^\omega$.

contains a club of $[H]^\omega$ or is disjoint from one, we conclude that $X(S) \cap H^\omega$ is either meager or comeager in H^ω and hence has the Baire Property in H^ω . The subspaces H^ω for $H \in C$ contain a club in our topological sense so we conclude that $X(S)$ has the Baire Property in κ^ω by $BRP(\kappa)$. If $X(S)$ is comeager in some neighborhood, then S contains a club, and if it is meager, then S is disjoint from a club. In either case, we have a contradiction. (3) is proved similarly. For (4) let $\mathcal{S} = \{S_\alpha \mid \alpha < \kappa\}$ be a sequence of stationary subsets of $[\kappa]^\omega$. For $x \in \kappa^\omega$ let \hat{x} be defined by $\hat{x}(n) = x(n+1)$. Define the space $X(\mathcal{S})$ to be the set of $x \in \kappa^\omega$ such that $\text{ran}(\hat{x}) \in S_{x(0)}$. Thus $X(\mathcal{S})$ is a Baire space by previous remarks. A club of $H \in [\kappa]^\omega$ at which \mathcal{S} does not reflect can be converted into a club of subspaces of κ^ω in which the trace of $X(\mathcal{S})$ is not Baire, and so (4) follows. \square

Theorem 1.2. *The following are equivalent.*

- (1) $BRP^w(\kappa) + WRP_{(2)}^c(\omega_2)$.
- (2) $BRP(\kappa)$.

Proof. Suppose $A \subseteq \kappa^\omega$ is such that $A \cap H^\omega$ has the Baire Property in H^ω for an ω_1 -club C of $H \in [\kappa]^\omega$. For $p \in \kappa^{<\omega}$ and $\sigma \in [\kappa]^\omega$ let $G_p^{**}(A, \sigma)$ denote the Banach-Mazur game where I and II alternate playing elements of $\sigma^{<\omega}$ which properly extend each other. I plays first and must extend p , and if $x \in \sigma^\omega$ is the cooperative play, then II wins if and only if $x \in A$. For a fixed $p \in \kappa^{<\omega}$ let S_i^p denote the set of $\sigma \in [\kappa]^\omega$ such that player i wins the Banach-Mazur game $G_p^{**}(A, \sigma)$, where i is either I or II . We first claim that $S_I^p \cup S_{II}^p$ contains a club for any $p \in \kappa^{<\omega}$. Suppose $S = [\omega_2]^\omega \setminus (S_I^p \cup S_{II}^p)$ is stationary. Then there is $H \in C$ such that $S \cap [H]^\omega$ is stationary by $WRP(\kappa)$. Since $A \cap H^\omega$ has the Baire Property in H^ω by assumption, there is a winning strategy τ in the game $G_p^{**}(A, H)$ for one of the players, without loss of generality player I , and hence for a club of $\sigma \in [H]^\omega$, the restriction $\tau \upharpoonright \sigma^{<\omega}$ witnesses $\sigma \in S_I^p$ giving the desired contradiction. We now show that S_I^p and S_{II}^p cannot both be stationary. Assume otherwise. By $WRP_{(2)}(\kappa)$ there is $H \in C$ such that both $S_I^p \cap H^\omega$ and $S_{II}^p \cap H^\omega$ are stationary in H^ω . Either I or II wins $G^{**}(A, H)$ via a strategy τ . Any σ closed under τ in the wrong set gives a contradiction. It now follows that for $p \in \kappa^{<\omega}$ either II wins $G_p^{**}(A, \sigma)$ for a club of σ or there is $p \subset q$ such that II wins $G_q^{**}(-A, \sigma)$ for a club of σ . To see this, fix p and suppose II fails to win $G_p^{**}(A, \sigma)$ for all σ in a stationary set S . By intersecting with a club and pressing down we find q so that II wins $G_q^{**}(-A, \sigma)$ for all σ in some stationary $\bar{S} \subseteq S$. If this set does not contain a club, then by repeating the previous argument we get an r extending q such that S_I^r and S_{II}^r are both stationary, contradicting the last claim. It now follows that A has the Baire Property. Suppose II wins $G_p^{**}(A, \sigma)$ on a club. Then II wins $G_p^{**}(A, H)$ for an ω_1 -club and so by $BRP^w(\kappa)$, II must win $G^{**}(A, \kappa)$. \square

As a corollary to Theorem 1.1, and the fact that $WRP(\omega_2)$ is equiconsistent with the existence of a weakly compact cardinal, we see that both $BRP^w(\omega_2)$ and $BRP(\omega_2)$ are as well. The global principle BRP^w is much stronger as it implies the global failure of \square_κ (see [13]). We now separate BRP^w from $BRP(\omega_2)$ using this connection with stationary reflection. The reader is referred to [10] for similar results in the context of MM .

Definition 1.3. \mathbb{P} consists of triples (p, e, δ) such that

- (1) $\delta < \omega_2$.
- (2) $e : \delta + 1 \times \omega_1 \rightarrow [\omega_2]^\omega$.
- (3) $(e(\gamma, \xi) \mid \xi < \omega_1)$ is continuous and exhaustive in $[\gamma]^\omega$ for each $\gamma \leq \delta$.
- (4) If $cf(\gamma) = \omega$, then $e(\gamma, \xi)$ is cofinal in γ for each $\xi < \omega_1$.
- (5) $p : ran(e) \rightarrow 2$.
- (6) For each $\gamma \leq \delta$ the set of $\sigma \in ran(e)$ such that $p(\sigma) = 0$ either contains or is disjoint from a club in $[\gamma]^\omega$.

The ordering is reverse inclusion on each coordinate. Note that clause (4) of the definition ensures that \mathbb{P} is σ -closed.

Lemma 1.4. \mathbb{P} is (ω_1, ∞) -distributive.

Proof. We show that II wins the game where I and II alternate playing a decreasing sequence of conditions $(p_\alpha, e_\alpha, \delta_\alpha)$ for $\alpha < \omega_1$ with II playing on the set of even stages E including limits. At the end, II must produce a p below every member of the sequence to win. From the existence of a winning strategy for II one easily deduces that \mathbb{P} is (ω_1, ∞) distributive (assuming I plays first). Fix $i < 2$. Note that II has complete control over e_γ at limit stages by clause (4). II wins by extending arbitrarily at successor stages, and at limits α setting

$$p_\alpha(e_\alpha(\delta_\alpha, \xi)) = i$$

for every $\xi < \omega_1$ and, via some simple bookkeeping, choosing the sequence $(e_\alpha(\delta_\alpha, \xi) \mid \xi < \omega_1)$ so that

$$\{e_\alpha(\delta_\alpha, 0) \mid \alpha \in E\}$$

is continuous and exhaustive in $[\delta]^\omega$ where δ is the limit of the δ_α . □

Lemma 1.5. Let $G \subset \mathbb{P}$ be V -generic. Suppose $WRP(\omega_2)$ holds in $V[G]$. Then $WRP_{(2)}^c(\omega_2)$ fails in $V[G]$.

Proof. The generic filter gives rise to $S^0, S^1 \subset [\omega_1]^\omega$ defined by

$$S^i = \bigcup \{p^{-1}(i) \mid \exists e, \delta (p, e, \delta) \in G\}$$

for $i < 2$ and a set

$$e = \bigcup \{ran(e) \mid \exists p, \delta (p, e, \delta) \in G\}.$$

A density argument shows that both S^0 and S^1 are stationary and that their union is equal to e . By design they do not simultaneously reflect. e contains a club in $[\omega_2]^\omega$ by $WRP(\omega_2)$, so the sets S_0, S_1 witness the failure of $WRP_{(2)}^c(\omega_2)$ in $V[G]$ as desired. □

The following basic result on extending embeddings to generic extensions is attributed to Silver (see [1] or Lemma 1 of [9]).

Lemma 1.6. Suppose M and N are inner models, $k : M \rightarrow N$ is elementary and $\mathbb{P} \in M$. Suppose \mathbb{P} is M -generic and $H \subset k(\mathbb{P})$ is N -generic. Suppose $k[G] \subset H$. Then there is a unique extension $k^* : M[G] \rightarrow N[H]$ of k with $k(G) = H$.

Theorem 1.7. BRP^w does not imply $BRP(\omega_2)$.

Proof. Let κ be supercompact and $G \subset \text{Col}(\omega_1, < \kappa)$ be V -generic. Let S_0, S_1 be $V[G]$ -generic subsets of $[\omega_2]^\omega$ which do not simultaneously reflect as given by the poset \mathbb{P} described above. If we can show that $BRP^w(\omega_2)$ holds in $V[G][S_0, S_1]$, then $BRP(\omega_2)$ must fail in this model by Theorem 1.1 and Lemma 1.5. In fact $BRP^w(\theta)$ holds for every θ , but for simplicity let us take $\theta = \omega_2$. Let $A \subset \omega_2^\omega$ have the property that $A \cap [H]^\omega$ is meager in H^ω for all H in some club $K \subseteq [\omega_2]^\omega$. Suppose toward a contradiction that A is of second category in ω_2^ω . Then one of the sets

$$A_0 = \{x \in A \mid \text{ran}(x) \in S^0\}$$

or

$$A_1 = \{x \in A \mid \text{ran}(x) \in S^1\},$$

say A_0 , is of second category. Let \mathbb{Q} be the poset which shoots a club through S_0 with conditions of size ω_1 . More specifically let p and e be the objects given by the generic \mathbb{P} and let D denote the set of $\delta < \omega_2$ such that $p(e(\delta, \xi)) = 0$ for every $\xi < \omega_1$. Let \mathbb{Q} be the poset of all continuous successor length sequences from D of size ω_1 ordered by reverse inclusion. If C is $V[G][S_0, S_1]$ -generic for \mathbb{Q} , then C gives rise to a club subset of S_0 . The idea is that the nonmeagerness of A_0 will be preserved in $V[G][S_0, S_1][C]$ where some generic supercompactness is recovered. Let (p, \dot{q}) be a condition in $\text{Col}(\omega_1, < \kappa) * \mathbb{P}$ and \dot{A} a term such that (p, \dot{q}) forces that \dot{A} is such a set as witnessed by \dot{K} and that \dot{A}_0 is nonmeager. Let $G^* \subset \text{Col}(\omega_1, < j(\kappa))$ be V -generic with $p \in G^*$ and let $G = G^* \cap \text{Col}(\omega_1, < \kappa)$. We thus have an embedding $j : V[G] \rightarrow M[G^*]$. In $V[G^*]$ the poset $\mathbb{P} * \mathbb{Q}$ has size ω_1 . Let $q = \dot{q}_G \in \mathbb{P}$. Build a condition $((S_0, S_1), C)$ in $\mathbb{P} * \mathbb{Q}$ which has a supremum in $j(\mathbb{P} * \mathbb{Q})$ and which meets every dense set in $V[G]$. Now let (S_0^*, S_1^*, C^*) be $M[G^*]$ -generic for $j(\mathbb{P} * \mathbb{Q})$ below this condition. By Lemma 1.6 above there is a unique extension

$$j^* : V[G][S_0, S_1][C] \rightarrow M[G^*][S_0^*, S_1^*][C^*].$$

Now we do the usual type of reflection and preservation argument. A_0 is nonmeager in $V[G][S_0, S_1][C]$. The factor

$$j(\mathbb{P} * \mathbb{Q}) / (S_0, S_1) * C$$

is σ -closed and hence $j^*[A] \subset j^*[A]^\omega$ is nonmeager in the space $j^*[\kappa]^\omega$ in the model $V[G^*][S_0^*, S_1^*][C^*]$ and hence in $M[G^*][S_0^*, S_1^*][C^*]$. It follows that the model $M[G^*][S_0^*, S_1^*][C^*]$ satisfies the sentence asserting the existence of $H \in [j(\kappa)]^{\omega_1}$ with $j^*(A) \cap H^\omega$ nonmeager in H^ω . By elementarity, $V[G][S_0, S_1][C]$ contains such an H . Thus $H \in V[G][S_0, S_1]$ as \mathbb{Q} does not add new sets of size ω_1 , and H must have the property there as well giving the desired contradiction. \square

Paul Larson noted that the above model can be modified so that $WRP_{(2)}(\omega_2)$ fails but $WRP_{(2)}^c(\omega_2)$ holds. Modify the poset \mathbb{P} so that disjoint stationary sets (S^0, S^1, S^2) are added whose union is $[\omega_2]^\omega$ and such that S^1 and S^2 do not simultaneously reflect. Suppose A and B are complimentary stationary sets. If A and B both have stationary intersection with some S^i , then design \mathbb{Q} to shoot a club through S^i and use the reflection argument above. Otherwise say A almost contains S^0 and B almost contains S^1 . Then A and B necessarily reflect simultaneously. Returning to Baire Reflection, we can deduce some corollaries. As a consequence of Theorem 1.2, $BRP_{(2)}^w(\omega_2)$ implies $WRP_{(2)}(\omega_2)$ and hence $BRP(\omega_2)$, and in the

model just described, $BRP(\omega_2)$ holds. Thus $BRP(\theta)$ does not imply $WRP_{(2)}(\theta)$ and hence does not imply $BRP_{(2)}^w(\theta)$.

2. BAIRE REFLECTION DOES NOT IMPLY CH

König shows in [9] that $GRP_\omega(\omega_2)$ implies CH . In this section we show that Baire Reflection is consistent with $2^\omega = \omega_2$. Let \mathbb{C}_κ be the poset for adding κ many Cohen reals. Conditions are finite partial functions from $\kappa \times \omega$ to 2 ordered by reverse inclusion.

Lemma 2.1. *Let κ and δ be arbitrary and suppose $G \subset \mathbb{C}_\kappa$ is V -generic. Suppose $A \subset \delta^\omega$ is nonmeager. Then A remains nonmeager in $V[G]$.*

Proof. Suppose $(\tau^n \mid n < \omega)$ is a sequence of nowhere dense sets such that

$$A \subseteq \bigcup_{n < \omega} \tau_G^n$$

in $V[G]$ for any $G \subset \mathbb{C}_\kappa$ which is V -generic. We define a Banach-Mazur strategy for player II in V as follows. Let θ be large enough and for $a \in H(\theta)$ let M_a be the skolem closure of a inside $H(\theta)$ together with a well-ordering of $H(\theta)$. We will always add the sequence (τ^n) and \mathbb{P} . Fix a bijection $n \rightarrow (n_0, n_1)$ from ω to $\omega \times \omega$. Given a sequence $\{s_0, s_1, \dots, s_n\}$ with each $s_i \in \delta^{<\omega}$ and $s_i \subseteq s_{i+1}$ for each i , let p be the n_1^{st} member of $M_{\{s_0, \dots, s_{n_0}\}} \cap \mathbb{P}$ and select q extending p and s extending s_n so that q forces that $[s]$ is disjoint from τ^i for each $i \leq n$, where $[s]$ is the basic open neighborhood of δ^ω determined by s . Set $s_{n+1} = s$. Since II cannot win the Banach-Mazur game for the compliment of A by this strategy there is a play

$$x = \bigcup \{s_n \mid n \in \omega\}$$

according to it which belongs to A . Let M be the skolem hull of the set of moves of this strategy, and M_n the hull of the first n moves. Thus $M = \bigcup_{n < \omega} M_n$. There is an $r \in \mathbb{P}$ such that r forces that $x \in \tau^m$ for some m . Let p be the restriction of r to M , and let k be big enough so that $k_0 = m$ and p is the k_1^{st} member of $M_{\{s_0, \dots, s_m\}}$. Then some extension of p forces that $[s_{m+1}]$ is disjoint from τ^m and hence that $x \notin \tau^m$. Such a condition can be found in $M_{\{s_0, \dots, s_{m+1}\}}$ so that it is compatible with r . This is a contradiction as $x \in [s_{m+1}]$. \square

We now briefly exposit (an equivalent version of) the model of [11]. For a set I let \mathbb{C}_I denote the poset for adding I many Cohen reals. Let $\dot{\sigma}_I$ denote the \mathbb{C}_I name for the set of all countable partial maps from I to 2. We work only with full names, that is, we assume that $\tau \in \dot{\sigma}_I$ is forced by \emptyset to be such a function. Let E be the set of even ordinals and O the set of odd ordinals. For an ordinal κ let $E_\kappa = E \cap \kappa$ and $O_\kappa = O \cap \kappa$.

Definition 2.2. \mathbb{M}_κ is the collection of all countable partial maps p such that

- (1) $\alpha \in E_\kappa \cap \text{dom}(p)$ implies $p(\alpha) \in 2^{<\omega}$,
- (2) $\alpha \in O_\kappa \cap \text{dom}(p)$ implies $p(\alpha) \in \dot{\sigma}_{E \cap \alpha}$,
- (3) $\{\alpha \in \text{dom}(p) \cap E_\kappa \mid p(\alpha) \text{ is nonempty}\}$ is finite.

\mathbb{M}_κ is ordered by setting $p \leq q$ if

- (1) $\text{dom}(q) \subseteq \text{dom}(p)$,
- (2) $\alpha \in \text{dom}(q) \cap E_\kappa$ implies $q(\alpha) \subseteq p(\alpha)$,
- (3) $\alpha \in \text{dom}(q) \cap O_\kappa$ implies $p \upharpoonright E_\alpha \Vdash_{\mathbb{C}_{E \cap \alpha}} q(\alpha) \subseteq p(\alpha)$.

Note the abuse of notation in (3). We are regarding $p \upharpoonright I$ as an element of \mathbb{C}_I for $I \subset E_\kappa$. The following lemma shows that preservation arguments in the context of the Mitchell collapse require only preservation under \mathbb{C}_{E_κ} and under σ -closed forcing.

Lemma 2.3. *There is a σ -closed poset \mathbb{P}_κ and a map $\pi : \mathbb{C}_{E_\kappa} \times \mathbb{P}_\kappa \rightarrow \mathbb{M}_\kappa$ such that whenever $G \subset \mathbb{C}_{E_\kappa} \times \mathbb{P}_\kappa$ is V -generic, then $\pi[G]$ is V -generic for \mathbb{M}_κ .*

Proof. \mathbb{P}_κ is the set of $p \in \mathbb{M}_\kappa$ which vanishes on even coordinates, that is, such that $\text{dom}(p) \cap E_\kappa$ is empty, with the ordering inherited from \mathbb{M}_κ . Clearly \mathbb{P}_κ is σ -closed. Define

$$\pi : \mathbb{C}_{E_\kappa} \times \mathbb{P}_\kappa \rightarrow \mathbb{M}_\kappa$$

by $\pi(a, p) = a \cup p$. Suppose $D \subset \mathbb{M}_\kappa$ is open and dense. Let $(a, p) \in \mathbb{C}_{E_\kappa} \times \mathbb{P}_\kappa$. Fix a maximal antichain of $b \in \mathbb{C}_{E_\kappa}$ below a such that there is $q_b \in \mathbb{P}_\kappa$ with $b \cup q_b \in D$ and $b \cup q_b \leq a \cup p$. The conditions q_b can be amalgamated (along with p) into a condition q such that each $b \cup q \in D$ and $q \leq p$ in \mathbb{P}_κ . \square

Theorem 2.4. *Suppose κ is supercompact and $G \subset \mathbb{M}_\kappa$ is V -generic. Then BRP holds in $V[G]$ but CH fails.*

Proof. See [10] for a proof that $2^\omega = \omega_2$ in $V[G]$. Fix λ and let $j : V \rightarrow M$ be an embedding with critical point κ such that $M^\theta \subset M$ for some large enough θ relative to λ . Let $G_{j(\kappa)} \subset \mathbb{M}_{j(\kappa)}$ be V -generic and $G_\kappa = G_{j(\kappa)} \cap \mathbb{M}_\kappa$. Thus j extends to

$$j : V[G_\kappa] \rightarrow M[G_{j(\kappa)}]$$

in $V[G_{j(\kappa)}]$. Since $WRP_{(2)}(\omega_2)$ holds in $V[G_\kappa]$ it suffices, by Theorem 1.2, to show that $BRP^\omega(\lambda)$ holds for any λ . So suppose $A \in V[G_\kappa]$ and $A \subset \lambda^\omega$ is not meager in λ^ω . We may assume that A is dense. Clearly $j[A] = j(A) \cap j[\lambda]^\omega$. Thus, if we can show that $j[A]$ is nonmeager in $j[\lambda]^\omega$ in the model $M[G_{j(\kappa)}]$, then $j[\lambda]$ is the desired element of $[j(\lambda)]^{\omega_1}$ and the result follows by reflection and the fact that $j[\lambda]$ is closed under $j(f)$ for any $f : \lambda^{<\omega} \rightarrow \lambda$ in $V[G_\kappa]$. Since j itself is a homeomorphism from λ^ω to $j[\lambda]^\omega$ which sends A to $j[A]$, $M[G_{j(\kappa)}] \subset V[G_{j(\kappa)}]$, and $j \upharpoonright \lambda^\omega \in V[G_{j(\kappa)}]$, it suffices to show that A is nonmeager in $V[G_{j(\kappa)}]$. The key point is that $V[G_{j(\kappa)}]$ is contained in an extension by a product of $\mathbb{C}_{E_{j(\kappa)}}$ and a σ -closed poset by Lemma 2.3. It is easy to see that the latter forcing preserves nonmeagerness, and the former is handled by Lemma 2.1. Thus A is nonmeager in an outer model of $V[G_{j(\kappa)}]$ and hence in $V[G_{j(\kappa)}]$ as desired. \square

As a final note, BRP does not imply $2^{\omega_1} = \omega_2$. The same is true of any consequence of GRP .

3. FAILURES OF BAIRE REFLECTION

Theorem 3.1. *Assume MM . Then every set of reals A of size ω_1 is ω_1 -Universally Baire but not ω_2 -Universally Baire.*

Proof. Suppose A is a set of reals of size ω_1 . Let $f : \omega_1^\omega \rightarrow \omega^\omega$ be continuous. Let O be the union of the set of open neighborhoods in ω_1^ω on which f is constant. If O is dense, then we are done. Let P denote the interior of $\omega_1^\omega \setminus O$. Thus P is nonempty and it suffices to show that $f^{-1}(A) \cap P$ has the Baire Property in P . Since P is homeomorphic to ω_1^ω we might as well assume that f is nowhere constant to begin with. Let B be the preimage of A and assume that B does not have the

Baire Property in ω_1^ω . Then there is a nonempty open set (again, which we take to be all of ω_1^ω) on which B and its complement are everywhere of second category. Consider the ideal

$$I = \{X \subset A \mid f^{-1}(X) \text{ is meager in } B\}.$$

I is a σ -ideal on a set of size ω_1 containing all singletons. We claim that I is ω_1 -dense in the sense that the quotient algebra $P(A)/I$ has a dense set of size ω_1 . This will be a contradiction by a result in [14] which shows that an ω_1 -dense ideal on a set of size ω_1 cannot exist under MA_{ω_1} . Consider an I -positive set X . By MA_{ω_1} (see [6]) we may write

$$X = \bigcup_{n < \omega} X_n,$$

where each X_n is closed in A . Thus

$$f^{-1}(X) = \bigcup_{n < \omega} f^{-1}(X_n),$$

where each $f^{-1}(X_n)$ is relatively closed in B . So there is an n such that the interior of $f^{-1}(X_n)$ is nonempty in the relative topology on B , which is to say, there is $s \in \omega_1^{<\omega}$ such that

$$\emptyset \neq [s] \cap B \subset f^{-1}(X_n),$$

where $[s] = \{x \in \omega_1^\omega \mid s \subseteq f\}$. Thus $f[[s] \cap B]$ is an I -positive subset of X . The claim follows since there are only ω_1 such sets.

We now show that A is not ω_2 -Universally Baire. It is more convenient to regard A as a subset of the unit interval $[0, 1]$. We may assume that A is dense. Let I be a precipitous ideal on A . For example, simply transfer the nonstationary ideal NS on ω_1 to A via a bijection between ω_1 and A . This ideal is precipitous (in fact saturated) under Martin's Maximum. This means that given any set $X \subset A$ with $X \notin I$ and sequence of I partitions $\{W_n \mid n \in \omega\}$ of X with W_{n+1} a refinement of W_n , there is a sequence $\{X_n \mid n \in \omega\}$ with nonempty intersection such that each X_n is an element of W_n (see [7, 8] for more details). Let \mathcal{B} be the set of open subintervals of $[0, 1]$ with rational endpoints. We define an operation on the power set of A by

$$\partial S = S \setminus \bigcup \{b \in \mathcal{B} \mid S \cap b \in I\}.$$

Note that $S \Delta \partial S \in I$ for any $S \subseteq A$. Define a metric space M as the set of pairs $(x, (X_n)) \in A \times I^\omega$ such that

- (1) $x \in \bigcap_{n < \omega} X_n$,
- (2) (X_n) is decreasing modulo I ,
- (3) there are infinitely many n such that $\partial X_n = X_n$.

M is a subspace of $[0, 1] \times I^\omega$ with the product topology and is therefore metrizable (I is discrete). The key use of the precipitous property is the following claim.

Claim 3.2. M is a Baire space.

Proof. Suppose that the empty player wins the Banach-Mazur game on M by a strategy τ . Here empty is the player who moves first and who wins if and only if the intersection of the neighborhoods played is empty. As usual, it suffices to restrict to basic open neighborhoods so we may assume that the output of τ is always such

a set. A typical basic open neighborhood in M is of the form $N_{b, (X_i)_{i \leq k}}$, for $b \in \mathcal{B}$ and each $X_i \subset b$, defined to be the set of $(x, (Y_n)) \in M$ such that

- (1) $x \in b \cap \bigcap_{i \in \omega} Y_i$.
- (2) $Y_i = X_i$ for every $i \leq k$.

Let \mathcal{N} be the set of open neighborhoods as above. For an element $N = N_{b, (X_i)_{i \leq k}} \in \mathcal{N}$ let ∂N denote $N_{b, (X_i)_{i \leq k+1}}$ where $X_{k+1} = \partial X_k$. Also let $l(N)$ denote X_k . Let N be the first move according to τ . We build a subtree $T \subset \mathcal{N}^{<\omega}$ as follows. The root of the tree is N . Suppose all sequences t of length $k+1$ have been defined. Given such a sequence $t = (N_0, N_1, \dots, N_k)$ suppose $N_k = N_{b, (X_i)_{i \leq j}}$. By varying τ we can produce a set of basic neighborhoods $\mathcal{A} \subset \mathcal{N}$ such that the set of $l(N)$ for $N \in \mathcal{A}$ is a maximal $P(A)/I$ antichain below $l(N_k)$. Sequences of the form (N_0, \dots, N_k, N) for $N \in \mathcal{A}$ will form the successors of t in T . For $t \in T$ let $X_t = l(N)$ where N is the last neighborhood on the sequence t . Let $W_k = \{X_t \mid t \in T\}$. Then the partitions W_k for $k \in \omega$ witness that the ideal I is not precipitous, giving the desired contradiction. \square

The projection map

$$\pi : M \rightarrow A,$$

which is continuous and nowhere constant, lifts to

$$\bar{\pi} : \beta M \rightarrow [0, 1].$$

The preimage $B = \bar{\pi}^{-1}(A)$ has the Baire Property in βM by assumption as βM has weight $2^{\omega_1} = \omega_2$. Thus B can be written as the symmetric difference of an open set U and a meager set F . As M is a Baire space M is of second category in βM and hence U is nonempty as $M \subseteq B$. Let $\{D_n \mid n \in \omega\}$ be open and dense subsets of U such that $\bigcap_{n \in \omega} D_n \subset B$. We can now build a Cantor scheme $\{U_s \mid s \in 2^{<\omega}\}$ of open subsets of U such that if $s \in 2^n$, then $U_s \subset D_n$ and for every s ,

$$\bar{\pi}[\bar{U}_{s0}] \cap \bar{\pi}[\bar{U}_{s1}] = \emptyset.$$

The set

$$P = \bigcap_n \bigcup_{s \in 2^n} \bar{U}_s$$

is perfect in βM by compactness and so $\bar{\pi}[P]$ is the desired perfect subset of A by design. \square

Corollary 3.3. *$BRP^w(\omega_2)$ fails under MM .*

Proof. Combine Theorem 3.1 with Theorem 1.2 and the fact that $WRP_{(2)}(\omega_2)$ holds under MM . \square

Theorem 3.4. *Assume MM . There is a Baire metric space of weight ω_2 with a club of subspaces of weight ω_1 which are meager in themselves. Hence $BRP^w(\omega_2)$ fails.*

Proof. For technical reasons we will work in the subspace $M \subset \omega_2^\omega$ of increasing sequences. M is a complete metric space homeomorphic to ω_2^ω . We will produce $X \subset M$ which is everywhere Baire but has the property that $X \cap \delta^\omega \cap M$ is meager in $\delta^\omega \cap M$ for every $\delta < \omega_2$. We will use MA_{ω_1} and $\diamond(S)$ where $S = \{\alpha < \omega_2 \mid cf(\alpha) = \omega\}$. Let $(d_\alpha \mid \alpha \in S)$ be a $\diamond(S)$ -sequence. We construct $X = \{x_\alpha \mid \alpha \in S\} \subset M$ as follows. If d_α codes a map $\pi_\alpha : \alpha^{<\omega} \rightarrow \alpha^{<\omega}$ and an increasing $p \in \omega_2^{<\omega}$ with

the property that $f(s)$ is increasing whenever s is increasing, then we view π_α as a Banach-Mazur strategy for player II in $\alpha^\omega \cap M$ and let $x_\alpha \in M$ be any legal play by this strategy where I has ensured that x_α is cofinal in α and extends p . Fix a comeager set $D \subseteq M$ and a neighborhood determined by a finite increasing sequence p . There is a winning strategy for II in the game associated to D , $d : \omega_2^{<\omega} \rightarrow \omega_2$, and there are club many $\alpha < \omega_2$ which are closed under this strategy. Thus d_α codes $d \upharpoonright \alpha^{<\omega}$ together with p for some $\alpha \in S$ and hence $x_\alpha \in D$ and extends p . This shows that X is a Baire space. We claim that $X \cap \delta^\omega \cap M$ is meager for every $\delta < \omega_2$. If $\delta \in S$ and $(\delta_n \mid n < \omega)$ is a sequence cofinal in δ , then

$$X \cap \delta^\omega = \{x_\delta\} \cup \bigcup_{n < \omega} X \cap \delta_n^\omega \cap M,$$

a countable union of sets meager in $\delta^\omega \cap M$. So let us assume δ has cofinality ω_1 . Here we use MA_{ω_1} . Let \mathbb{P} be the set of conditions

$$p = (\Gamma_p, X_p)$$

such that Γ_p is a finite subset of δ and X_p is a finite subset of $S \cap \delta$. For a pair of conditions p, q we define $p \leq q$ if $\Gamma_q \subseteq \Gamma_p$, $X_q \subseteq X_p$ and for every $\alpha \in X_q$ we have $(\Gamma_p \setminus \Gamma_q) \cap \text{ran}(x_\alpha) = \emptyset$.

Claim 3.5. \mathbb{P} satisfies the countable chain condition.

Proof. Suppose $(p_\alpha \mid \alpha < \omega_1)$ is an antichain. By two applications of the Δ -system lemma we may assume that there are sets s and t such that

$$\Gamma_{p_\alpha} \cap \Gamma_{p_\beta} = s \text{ and } X_{p_\alpha} \cap X_{p_\beta} = t$$

for every $\alpha < \beta < \omega_1$. There is a subsequence which we reindex as

$$(p_\alpha \mid \alpha < \omega^2 + 1)$$

such that for every $\alpha < \beta < \omega^2$,

$$\max(\Gamma_{p_\alpha} \setminus s) < \min(X_{p_\beta} \setminus t).$$

It now follows that some x_γ with $\gamma \in X_{p_{\omega^2+1}}$ has order type greater than ω which is a contradiction. \square

If $G \subset \mathbb{P}$ is a filter and G meets the requisite dense sets, then

$$\Gamma = \bigcup \{ \Gamma_p \mid p \in G \}$$

is an unbounded subset of δ which is almost disjoint from x_α for every $\alpha \in S \cap \delta$. An easy argument shows that each

$$X_k = \{x_\alpha \mid \alpha \in S \cap \delta \text{ and } \forall n > k \ x_\alpha(n) \in \delta \setminus \Gamma\}$$

is meager, from which it follows that $X \cap \delta^\omega \cap M$ is meager in $\delta^\omega \cap M$. \square

Theorem 3.6. *$BRP^w(2^\omega)$ for point countable spaces fails under MA_{ω_1} .*

Proof. We will use the tree $\sigma\mathbb{Q}$ of countable well-ordered increasing sequences of rationals which are bounded. We do not require the sequences to be continuous. For $t \in \sigma\mathbb{Q}$ and $q \in \mathbb{Q}$ let $B_{t,q}$ denote

$$\{s \in \sigma\mathbb{Q} \mid t \subseteq s \text{ and } \sup(s) < q\}.$$

These sets together with their complements form a point countable base for a zero-dimensional topology on $\sigma\mathbb{Q}$. We claim that $\sigma\mathbb{Q}$ is a Baire space. In the

relevant Banach-Mazur game, the nonempty player wins by ensuring that the t_n are increasing, where $\{B_{t_n, q_n} \mid n < \omega\}$ is the sequence of plays, in which case

$$\bigcup_{n < \omega} s_n \in \bigcap_{n < \omega} B_{t_n, q_n}.$$

An ω_1 -club of closed subspaces of $\sigma\mathbb{Q}$ are subtrees of $\sigma\mathbb{Q}$ (viewed as a tree in the obvious way), and have no isolated points. Simply close under the successor and predecessor function. Any such T has size ω_1 and hence is special under MA_{ω_1} (see [16] for example). Let T be such a subspace and $\pi : T \rightarrow \omega$ a specializing map. Each $\pi^{-1}(n)$ is closed and nowhere dense in T . Thus T is meager in itself. \square

We conclude with some observations concerning reflection of second category to subspaces of countable weight. The principle analogous to $BRP^\omega(\kappa)$ would assert that $X \subset \kappa^\omega$ is meager iff $X \cap \sigma^\omega$ is meager for a club of $\sigma \in [\kappa]^\omega$. In the case $\kappa = \omega_1$ a counterexample exists under \diamond , and we suspect that the principle is true under the Proper Forcing Axiom, though we do not have a proof at present. For $\kappa = \omega_2$, the principle is outright false.

Theorem 3.7. *There is $X \subset \omega_2^\omega$ of second category such that $X \cap \sigma^\omega$ is meager in σ^ω for a club of $\sigma \in [\omega_2]^\omega$.*

Proof. Fix a sequence $\{C_\alpha \mid \alpha < \omega_1 \text{ and } cf(\alpha) = \omega\}$ with each $C_\alpha : \omega \rightarrow \alpha$ cofinal and increasing, having the property that every club subset of ω_2 includes the range of one (and hence stationarily many) C_α (see [12]). Fix a surjective map $\phi : [\omega]^\omega \rightarrow [\omega_1]^\omega$. A real $r \in 2^\omega$ codes a sequence of reals $\{r_n \mid n < \omega\}$ defined by $r_n(k) = r(2^n(2k+1))$. We associate another operation $\phi^* : [\omega]^\omega \rightarrow [\omega_1]^\omega$ defined by

$$\phi^*(r) = \bigcup_{n < \omega} \phi[r_n],$$

where we identify a real $s \in 2^\omega$ with $s^{-1}(1) \in [\omega]^\omega$. For $\sigma \in [\omega_2]^\omega$ define ν_σ by

$$\nu_\sigma = \pi_\sigma^{-1}(\phi^*(osc(\sigma)))$$

where $\pi_\sigma : \sigma \rightarrow otp(\sigma)$ is the collapse of σ and $osc(\sigma)$ is defined to be the set

$$\{n \mid \sigma \cap [C_{sup(\sigma)}(n), C_{sup(\sigma)}(n)) \neq \emptyset\}.$$

The sequence $\{\nu_\sigma \mid \sigma \in [\omega_2]^\omega\}$ can be shown to have the following property: for any $X \subset \omega_2$ the set of $\sigma \in [\omega_2]^\omega$ such that $X \cap \sigma = \nu_\sigma$ is stationary in $[\omega_2]^\omega$ (see Lemma 43 of [15] and [5]). We may view ν_σ as coding a comeager subset of σ^ω (as in Theorem 3.4) in some canonical way. For example, fix a bijection

$$c : \omega_2 \times \omega_2^{<\omega} \rightarrow \omega_2,$$

and observe that if $Y \subset \omega \times \omega_2^{<\omega}$ is a comeager set in the obvious way ($\hat{Y} = \{x \in \omega_2^\omega \mid \forall n \exists k (n, x \upharpoonright k) \in Y\}$ is comeager), then $c^{-1}(c(Y) \cap \sigma) = \hat{Y} \cap \omega \times \sigma^{<\omega}$ is comeager in σ^ω for a club of $\sigma \in [\omega_2]^\omega$. Define the space $X \subset \omega_2^\omega$ to be $\{x_\sigma \mid \sigma \in [\omega_2]^\omega \text{ where } x_\sigma \text{ is an element of } c^{-1}(\nu_\sigma) \text{ in the case that this is a code for a comeager set in } \sigma^\omega, \text{ and such that the range of } x_\sigma = \sigma. \text{ It is clear that } X \text{ is of second category as for any comeager set } Y \subset \omega_2^\omega \text{ there is } \sigma \text{ such that } c^{-1}(\nu_\sigma) = Y \cap (\omega \times \sigma^{<\omega}) \text{ so that } x_\sigma \text{ is in } \hat{Y}. \text{ On the other hand, } X \cap \sigma^\omega \text{ is a singleton (or empty), and hence meager, for every } \sigma \in [\omega_2]^\omega. \square$

REFERENCES

- [1] Cummings, J., *Large Cardinal Properties of Small Cardinals*, 1988, unpublished manuscript
- [2] Engelking, R., *General Topology*, Heldermann Verlag, 1989 MR1039321 (91c:54001)
- [3] Feng, Q., Jech, T., *Local Clubs, Reflection, and Preserving Stationary Sets*, Proceedings of the London Mathematical Society, Vol. 58, 1989, 237-257 MR977476 (90a:03072)
- [4] Feng, Q., Magidor, M., Woodin, H., *Universally Baire Sets of Reals*, MSRI Publications 26, 1992 MR1233821 (94g:03095)
- [5] Foreman, M., Todorcevic, S., *A New Löwenheim-Skolem Theorem*, Transactions of the American Mathematical Society 357, 2005, 1693-1715 MR2115072 (2005m:03064)
- [6] Fremlin, D.H., *Consequences of Martin's Axiom*, Cambridge University Press, 1984 MR780933 (86i:03001)
- [7] Galvin, F., Jech, T., Magidor, M., *An Ideal Game*, Journal of Symbolic Logic, Vol. 43, No. 2, 1978, 284-292 MR0485391 (58:5237)
- [8] Jech, T., Magidor, M., Mitchell, M., Prikry, K., *Precipitous Ideals*, Journal of Symbolic Logic, Vol. 45, No. 1, 1980, 1-8 MR560220 (81h:03097)
- [9] König, B., *Generic Compactness Reformulated*, Archive for Mathematical Logic 43, 2004, no. 3, 311-326 MR2052885 (2005f:03079)
- [10] Larson, P., *Separating Stationary Reflection Principles*, Journal of Symbolic Logic, Vol. 65, No. 1, 2000, 247-258 MR1782117 (2001k:03094)
- [11] Mitchell, W., *Aronsojn Trees and the Independence of the Transfer Property*, Annals of Mathematical Logic, Vol. 5, No. 1, 1975, 21-46 MR0313057 (47:1612)
- [12] Shelah, S., *Cardinal Arithmetic*, Vol. 29 of Oxford Logic Guides, Oxford University Press, 1994 MR1318912 (96e:03001)
- [13] Steel, J., *PFA implies $AD^{L(\mathbb{R})}$* , Journal of Symbolic Logic, Vol. 45, No. 4, 2005, 1255-1296 MR2194247
- [14] Taylor, A.D., *Regularity Properties of Ideals and Ultrafilters*, Annals of Mathematical Logic 16, 1979, No. 1, 33-55 MR530430 (83b:04003)
- [15] Todorcevic, S., *Localized Reflection and Fragments of PFA*, DIMACS Series in Discrete Mathematics and Theoretical Computer Science, vol. 58, 2002, 135-148 MR1903856 (2003g:03081)
- [16] Todorcevic, S., *Trees and Linearly Ordered Sets*, Handbook of Set Theoretic Topology, North Holland, 1984, 235-293 MR776625 (86h:54040)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, CANADA M5S 2E4 –
 AND – UNIVERSITE PARIS 7-CNRS, UMR 7056, 2 PLACE JUSSIEU, 75251 PARIS CEDEX 05, FRANCE
E-mail address: stevo@math.toronto.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, CANADA M5S 2E4
Current address: Department of Mathematics, Wesleyan University, 265 Church Street, Middletown, Connecticut 06459-0128
E-mail address: azoble@wesleyan.edu